



**NO** compromises

**TAZOREN**<sup>TM</sup>  
Piperacillin and Tazobactam for Inj. 4.5 gm

la Renon<sup>®</sup>

## No Compromises

### Indications:

**TAZOREN™** is indicated for treatment of the following systemic and/or local bacterial infections:

#### **Adults and the Elderly**

- Lower respiratory tract infections.
- Urinary tract infections. (complicated and uncomplicated)
- Intra-abdominal infections including peritonitis and appendicitis complicated with rupture or abscess.
- Skin and skin infections. (complicated and uncomplicated)
- Post-partum endometritis and pelvic infections.
- Polymicrobial infections.
- Bacterial septicaemia.
- Bacterial infections in neutropenic adults in combination with an aminoglycoside.

#### **Pediatrics**

- Intra-abdominal infections including appendicitis complicated by rupture with peritonitis and/or abscess formation, biliary infections.
- Bacterial infections in neutropenic children in combination with an aminoglycoside.

### Composition:

Each vial contains

Piperacillin sodium equivalent to Piperacillin..... 4gm

Tazobactam sodium equivalent to Tazobactam..... 500 mg

### Dosage and Administration:

**TAZOREN™:** Slow intravenous infusion (e.g., over 20-30 minutes) or slow intravenous injection (over at least 3-5 minutes).

#### **Adults and Children 12 Years and Older:**

**General Infections:** 12 g piperacillin /1.5 g tazobactam/day in divided doses every 6 or 8 hours.

**Severe Infections:** 18g piperacillin /2.25 g tazobactam/day in divided doses.

#### **Pediatric Intra-Abdominal Infection:**

**Children (2 to 12 years, >40kg and normal renal function):** 100 mg piperacillin/12.5mg tazobactam/Kg BW /8hours.

### Storage Conditions:

Injection vials should be stored at (20<sup>0</sup>-25°C) room temperature prior to reconstitution.