


*Navigate the Road of Life Ahead*



In Bipolar Disorder & Scizophrenia,

*Recommend,*


**ARIPIREN™**

**Aripiprazole 5 mg / 10 mg / 15 mg / 20 mg Tablets**

**La Renon®**

In Bipolar Disorder & Scizophrenia,

Recommend.  
**ARIPIREN™**  
 Aripiprazole 5 mg / 10 mg / 15 mg / 20 mg Tablets



**Indication**

**ARIPIREN™** is indicated for –

- A. Schizophrenia
- B. Acute treatment of Manic and Mixed episodes associated with Bipolar I Disorder
- C. Adjunctive treatment of Major Depressive Disorder
- D. Irritability associated with Autistic Disorder
- E. Treatment of Tourette's Disorder

**Mechanism of action**

Aripiprazole's antipsychotic activity is likely due to a combination of antagonism at D2 receptors in the mesolimbic pathway and 5HT2A receptors in the frontal cortex. Antagonism at D2 receptors relieves positive symptoms while antagonism at 5HT2A receptors relieves negative symptoms.

**Dosage**

| Indication   | Initial Dose | Recommended Dose | Maximum Dose |
|--|--------------|------------------|--------------|
| <b>Schizophrenia</b>   |              |                  |              |
| Adults   | 10-15 mg/day | 10-15 mg/day     | 30 mg/day    |
| <b>Bipolar Disorder- Treatment of manic or mixed episodes</b>      |              |                  |              |
| Bipolar Mania Adults Monotherapy                                   | 15 mg/day    | 15 mg/day        | 30 mg/day    |
| Bipolar Mania Adults Adjunct to lithium or valproate               | 10-15 mg/day | 15 mg/day        | 30 mg/day    |
| <b>Major Depressive Disorder</b>                                   |              |                  |              |
| Adjunct to antidepressants for the treatment of MDD                | 2-5 mg/day   | 5-10 mg/day      | 15 mg/day    |
| <b>Autism Spectrum Disorders</b>                                   |              |                  |              |
| Irritability associated with autistic disorder. Pediatric patients | 2 mg/day     | 5-10 mg/day      | 15 mg/day    |

**Advantage**

- A. Aripiprazole is effective in reducing risperidone-induced hyperprolactinemia and its sequelae\*.
- B. Clinically stable patients with schizophrenia who are taking olanzapine and who have evidence of metabolic syndrome can be successfully switched to aripiprazole, with improvement in various parameters of metabolic syndrome and without any significant change in efficacy measures.\*\*

**Food Interactions:**

None. No effect with intake of food.

\*Neuropsychiatric Disease and Treatment 2015:11 549-555

\*\* Neuropsychiatric Disease and Treatment 2015:11 685-693



**La Renon Healthcare**

207-208 Iscon Elegance, Circle P, Prahlad Nagar Cross Roads  
 S.G. Highway, Ahmedabad-380015, Gujarat, India.  
 Phone: + 91-79-3046-1000 (30 lines) Fax: +91-79-3046-1001  
 E-mail: info@larenon.com Web: www.larenon.com

I am \_\_\_\_\_  
 Call me on \_\_\_\_\_  
 Mail me at \_\_\_\_\_