

LAFORMIN G-1/G-2

Glimepiride 1 mg/2 mg + Metformin (SR) 500 mg Tablets

LAFORMIN 6-1 FORTE/6-2 FORTE

Glimepiride 1 mg/2 mg + Metformin (SR) 1000 mg Tablets

<u>La Renon</u>

LAFORMIN G-1/G-2

Glimepiride 1 mg/2 mg + Metformin (SR) 500 mg Tablets

LAFORMIN G-1 FORTE/G-2 FORTE

Glimepiride 1 mg/2 mg + Metformin (SR) 1000 mg Tablets

Background:

- According to ADA & EASD consensus guidelines, many patients can be managed effectively with monotherapy, but disease
 is progressive. So they will require use of combination therapy in many, if not in most patients over time to maintain
 glycaemia in target range.
- When adding second drug, synergy of particular combination and other interactions should be considered. Antihyperglycaemic agents with different mechanisms of action will have greatest synergy.

Description:

Laformin G-1/G-2 and G-1 Forte/G-2 Forte uncoated bilayer Tablet is Oral Anti hyperglycemic tablet containing Glimepiride and Metformin used in the management of Type 2 Diabetes Mellitus and in particular, in Overweight and Obese people.

Indication:

Laformin G-1/G-2 and G-1 Forte/G-2 Forte is indicated as an adjunct to diet and exercise to improve glycemic control in patients with type-2 diabetes whose diabetes is not adequately controlled with metformin alone, or for those patients who have initially responded to glimepiride alone and require additional glycemic control.

Key Features of Laformin G-1/G-2 and G1 Forte/G2 Forte

- > Glimepiride and Metformin act complementary to each other.
- > Both compounds have a synergistic antihyperglycaemic effect without increasing the adverse effects of either pharmacological class
- > It provide Dosage Flexibility, hence Better compliance.
- > It do not cause Hypoglycemia
- > It is associated with less weight gain compared to other drug therapies.

Dosage:

- Laformin G-1/G-2 and G-1 Forte/G-2 Forte should be given once daily with meals and should be started at a low dose. Dosage should be individualized on the basis of both effectiveness and tolerance.
- The initial recommended dose is one tablet once daily with breakfast or first main meal of the day.

Presentation:

Laformin G-1/G-2 and G-1 Forte/G-2 Forte is available as a strip of 10 Tablets.